arXiv:math/0403065v1 [math.DG] 3 Mar 2004

EIGEN-VALUE MONOTONICITY FOR THE RICCI-HAMILTON FLOW

LI MA

Abstract. In this short note, we discuss the monotonicity of the first eigenvalue of the Laplacian operator to the Ricci-Hamilton flow on a compact or a complete non-compact Riemannian manifold.

Let (M, g_0) be a compact or a complete non-compact Riemannian manifold. We study the Ricci-Hamilton flow of a family of Riemannian metrics $\{g(t)\}$ with $g(0) = g_0, 0 \leq t < T$. Let $g = g(t)$. In the local coordinates (x^i), we write

$$g = g_{ij} dx^i dx^j.$$

By definition, the Ricci-Hamilton flow is defined by the evolution equation:

$$\partial_t g_{ij} = -2R_{ij}, \quad \text{on } M_T := M \times [0, T)$$

where R_{ij} is the Ricci tensor of the metric $g := g(t)$ and T is the maximal existing time for the flow. In [2], R.Hamilton proved the local existence of the flow for the compact manifold case. His argument is much simplified by De Turck [1]. When (M, g_0) is a a complete non-compact Riemannian manifold with bounded geometry, W.X.Shi [5] obtained the local existence result for the flow.

We study the monotonicity property of the first eigenvalue of the Laplacian operator $\Delta := \Delta_{g(t)}$ of the evolving metric $(g(t))$ along the Ricci-Hamilton flow. More precisely, let D be a compact domain with smooth boundary in the manifold M. Let $f := f(x, t)$ be the first eigen-function of Δ. Then we have

$$-\Delta f = \mu f, \quad \text{in } D, \tag{1}$$

with Dirichlet boundary condition

$$f = 0, \quad \text{on } \partial D$$

where $\mu := \lambda_1(D) > 0$ is the corresponding first eigen-value of Δ. We can normalize f such that

$$\int_D f^2 dv = 1. \tag{2}$$

Here dv is the volume form of the metric $g = g(t)$.

We will use the following conventions: For $F = F(t)$ being a quantity on D, we let F' be the derivative with respect to the time variable t and

---

1991 *Mathematics Subject Classification.* 53C44.

Key words and phrases. Ricci-Hamilton flow, eigen-value.

let $F_i, F_{ij}, F_{ij;k}$, etc, be the covariant derivatives of F with respect to the Levi-Civita connection of the metric g. We also assume that when $D = M$ is compact, we impose the natural condition that $\int_M f dv = 0$. We abuse the upper or lower indexes in this paper when their meanings are clear. We write D_T for $D \times [0, T)$.

Differentiating (1), we get

$$-\Delta' f - \Delta f' = \mu' f + \mu f'. \quad \text{on } D, \tag{3}$$

Multiplying both sides of (3) by f and integrating over D, we have

$$-\int \Delta' f \cdot f - \int \Delta f' \cdot f = \mu' + \mu \int f' f. \tag{4}$$

Note that

$$-f' \Delta f = \mu f' f, \quad \text{on } D.$$

Then we have

$$\mu' = -\int \Delta' f f$$

Hence using (7) in the appendix we obtain that

$$\mu' = 2 \int R_{ij} f_{ij} f.$$

If $n = 2$, then we have

$$R_{ij} = \frac{1}{2} R g_{ij},$$

and

$$\mu' = \int R \Delta f \cdot f = -\mu \int f^2 R dv \leq 0,$$

provided $R > 0$.

In the case when M is a closed surface and $R \geq C > 0$ for some uniform constant C, we have

$$(\log \mu)' \leq -C.$$

It is also clear that when $R < 0$, we have

$$\mu' = \int R \Delta f \cdot f = -\mu \int f^2 R dv > 0.$$

When $n \geq 3$, we notice that

$$\begin{aligned}
\int R_{ij} f_{ij} f &= -\int (R_{ij} f)_j f_i \\
&= -\int R_{ij;j} f f_i - \int R_{ij} f_i f_j \\
&= -\frac{1}{2}\int R_i f_i f - \int R_{ij} f_i f_j \\
&= \frac{1}{2}\int R (f_i f)_i - \int R_{ij} f_i f_j \\
&= \frac{1}{2}\int R \Delta f f + \frac{1}{2}\int R |\nabla f|^2 - \int R_{ij} f_i f_j \\
&= \frac{-\mu}{2}\int R f^2 - \int (R_{ij} - \frac{R}{2} g_{ij}) f_i f_j.
\end{aligned}$$

Here we have used the contracted second Bianchi identity

$$2R_{ij;j} = R_i.$$

Therefore, we have proved the following result:

Main Theorem . *Let $g = g(t)$ be the evolving metric along the Ricci-Hamilton flow with $g(0) = g_0$ being the initial metric. Let D be a smooth bounded domain in (M, g_0). Let $\mu > 0$ be the first eigen-value of the Laplacian operator of the metric $g(t)$. If the scalar curvature $R \geq 0$ and non-trivial in D_T and the Einstein tensor*

$$E_{ij} := R_{ij} - \frac{R}{2} g_{ij} \geq 0, \quad \text{in } D_T,$$

then we have $\mu' < 0$, that is to say, μ is strictly decreasing along the Rici-Hamilton flow. On the contrary, if the scalar curvature $R \leq 0$ in D_T and the Einstein tensor

$$E_{ij} \leq 0, \quad \text{in } D_T,$$

then we have $\mu' \geq 0$.

1. APPENDIX

Given a Riemannian manifold (M^n, g). In the local coordinates (x^i), we write

$$g = g_{ij} dx^i dx^j.$$

Let u be a smooth function. Then the Laplacian of u is defined by

$$\Delta u = \frac{1}{\sqrt{|g|}} \partial_i (\sqrt{|g|} g^{ij} \partial_j u)$$

where

$$(g^{ij}) = (g_{ij})^{-1}$$

is the inverse of the matrix (g_{ij}) and $|g| = det(g_{ij})$.

For convenient of readers, we review some facts about Ricci-Hamilton flow. Recall that the metric $g = g(t)$ satisfies the Ricci-Hamilton flow:

$$\partial_t g_{ij} = -2R_{ij}, \quad \text{on } M_T.$$

Along this flow, we have

$$\partial_t dv = -R dv, \tag{5}$$

where dv is the volume element and R is the scalar curvature of the metric $g(t)$ respectively. For smooth functions u and v with compact support, we have by a use of the divergence theorem that

$$\int g^{ij} u_i v_j dv = -\int \Delta u v dv.$$

Differentiating both sides of this equation and using (5) we have

$$2\int R_{ij}u_iv_j - \int Ru_iv_i = -\int \Delta' uv + \int \Delta u \cdot vR. \tag{6}$$

Since

$$\int Ru_iv_i = \int (Ru_i)_i v = \int R_iu_iv + \int Rv\Delta u,$$

we have that

$$2\int R_{ij}u_iv_j + \int (\nabla R, \nabla u)\, v = \int \Delta' uv.$$

As before, using integration by part and using the contracted second Bianchi identity, we obtain that

$$\int R_{ij}u_iv_j = -\frac{1}{2}\int (\nabla R, \nabla u)\, v - \int R_{ij}u_{ij}v.$$

Combining this with (6) we get

$$\int \Delta' uv = -2\int R_{ij}u_{ij}v.$$

Hence we have the variation formula for the Laplacian operator to the Ricci-Hamilton flow:

$$\Delta' u = -2R_{ij}u_{ij}. \tag{7}$$

In particular when $n = 2$, we have

$$R_{ij} = \frac{1}{2}Rg_{ij}.$$

For more material about Ricci-Hamilton flow, one may see [4] and [3].

References

[1] De Turck, *Deforming metrics in the direction of their Ricci tensors*, J.Diff. Geom., 18(1983)157-162.

[2] R.Hamilton, *The Ricci flow in dimension three*, J. Diff. Geom., 17(1982)255-306.

[3] Li Ma, *Ricci-Hamilton flow on surfaces*, (a book to be published by Global Science Publishing), Singapore, 2004.

[4] G.Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math.DG/0211159.

[5] W.X.Shi, *Deforming the metric on complete Riemannian manifolds*, J. Diff. Geom. 30(1989)353-360.

MA: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA

E-mail address: `lma@math.tsinghua.edu.cn`